

Field Equations of the $CP(N-1)$ Affine Gauge Theory

P.Leifer^{1a} and L.P.Horwitz^{abc}

^aBar-Ilan University, Ramat-Gan, Israel;

^bSchool of Physics and Astronomy, Tel-Aviv University, Ramat Aviv, Israel;

^cCollege of Judea and Shomron, Ariel, Israel

Abstract

A non-linear relativistic 4D field model of a quantum particle which emerges from the internal dynamics in the quantum phase space $CP(N-1)$ is proposed. In this model there is no distinction between ‘particle’ and its ‘surrounding field’, and the space-time manifold emerges from the description of the quantum state. The quantum observables of the ‘quantum particle field’ are described in terms of the affine parallel transport of the local dynamical variables in $CP(N-1)$.

PACS numbers: 03.65.Ca, 03.65.Ta, 04.20.Cv

1 Introduction

We discuss in this paper a simple model of affine gauge field equations describing some generalized coherent state. This model admits both Goldstone and Higgs modes of evolution, whose dynamical properties may be described in terms of nonlinear relativistic wave equations. The model, furthermore, has the property that the physical particle includes, by definition, its self-interaction in a natural way. The basic structure of the model, involving the natural motions in a $CP(N-1)$ manifold induced by the generators of $SU(N)$ has been discussed elsewhere [1, 2, 3]. The invariant Cartan decomposition of the Lie algebra $AlgSU(N)$ of generators expressed in terms of the local coordinates admits an invariant embedding of the isotropy sub-group $H = U(1) \times U(N-1) \subset G = SU(N)$ and the coset transformations $G/H = SU(N)/[U(1) \times U(N-1)] = CP(N-1)$ of the generalized coherent state (GCS). The symplectic and Riemannian structure of the $CP(N-1)$ manifold may be expressed in local coordinates as well [4, 5, 6, 7]. The normal and tangent vectors at a point on

¹On leave from Crimea State Engineering and Pedagogical University, Simferopol, Crimea, Ukraine

the $CP(N-1)$ manifold form the unitary basis of a spinor corresponding to an observable associated with a two state system, as we shall explain below. The affine connection associated with the Fubini-Study metric gives an invariant relation between such vectors at nearby points which can be described in terms of the transformations of $SL(2, C)$. There is a close analogy between those and Lorentz transformations in the actual space-time. In this framework, we construct nonlinear relativistic field equations which are covariant under the diffeomorphisms generated by the local actions of $SU(N)$, realized in terms of the $SL(2, C)$ action on the local spinor in the state space C^2 . We review the basic structure briefly in the following, and show how the field equations arise. We finally discuss some properties of the solutions of these equations which could be associated with elementary particles.

The main new points of our approach are as follows:

A. We use the notion of “elementary quantum motion states” with well defined quantized Planck’s action $S_a = \hbar a$. Their GCS serve as an abstract formalization of the “quasi-classical” description of a quantum setup or “Schrödinger’s lump” [8].

B. The quantum phase space $CP(N-1)$ serves as the base of a fibre bundle for which the tangent space corresponds to local dynamical variables (LDV’s). The particular section of this bundle and the corresponding affine gauge field are geometric tools for the construction of a theory admitting quantum measurement in the state-dependent dynamical space-time.

The technical details are as follows:

1. The projective representation of pure N -dimension quantum states (one could think of arbitrary large N), provides a natural non-linear realization of the $G = SU(N)$ group manifold and the coset sub-manifold $G/H = SU(N)/S[U(1) \times U(N-1)] = CP(N-1)$. We shall consider the generators of this group as local dynamical variables [3] of the model.

2. These quantum dynamical variables are represented by the tangent vector fields to $CP(N-1)$. Embedding of $CP(N-1)$ into $\mathcal{H} = C^N$ provides a measurement procedure for the dynamical variables.

3. This notion of measurement gives rise to the analog of a local dynamical space-time capable of detecting the coincidence of expectation and measured values of these quantum dynamical variables.

4. The affine parallel transport, associated with the Fubini-Study metric, accompanied with “Lorentz spin transformation matrix” [9], establish this coincidence due to the identification of the parallel transported LDV in different GCS’s.

5. The parametrization of the measurement, with the help of *attributed* local space-time coordinates, is in fact an embedding of quantum dynamics in Hilbert space into a 4D world. This procedure is well defined due to the existence of the infinitesimal $SL(2, C)$ transformations of the spinor treated as Lorentz transformations of local space-time coordinates.

Below are introduced some fundamental notions of our construction.

2 The Action Quantization

We discuss a modification of the “second quantization” procedure.

First. In the second quantization method one has formally given particles whose properties are defined by some commutation relations between creation-annihilation operators. Note, that the commutation relations are only the simplest consequence of the curvature of the dynamical group manifold in the vicinity of the group’s unit (in algebra). Dynamical processes require, however, finite group transformations and, hence, the global group structure. The main technical idea is to use vector fields over a group manifold instead of Dirac’s q-numbers. This scheme therefore seeks the dynamical nature of the creation and annihilation processes of quantum particles.

Second. We shall primarily quantize the action, and not the energy. The relative (local) vacuum of some problem is not necessarily the state with minimal energy, it is a state with an extremal of some action functional.

POSTULATE 1.

We assume that there are elementary quantum states $|\hbar a\rangle, a = 0, 1, \dots$ of an abstract Planck oscillator whose states correspond to the quantum motions with given number of Planck action quanta.

We shall construct non-linear field equations describing energy (frequency) distribution, whose soliton-like solution provides the quantization of the dynamical variables. Quantum “particles”, and, hence, their numbers arise as some countable solutions of non-linear wave equations. In order to establish acceptable field equations which are capable of intrinsically describing all possible degrees of freedom under intensive interaction we construct a *universal ambient Hilbert state space* $\mathcal{H}$. We will use *the universality of the action* whose variation is capable of generating any relevant dynamical variable. We shall call vectors of the action state space $\mathcal{H}$ *action amplitudes*. Some of them will be elementary quantum states of motion corresponding to discrete numbers of Planck’s quanta $|\hbar a\rangle$. The action may create a linear superposition of $|\hbar a\rangle = (a!)^{-1/2}(\hat{\eta}^+)^a|0\rangle$ constituting $SU(\infty)$ multiplete of the Planck’s action quanta operator $\hat{S} = \hbar\hat{\eta}^+\hat{\eta}$ with the spectrum $S_a = \hbar a$ in the separable Hilbert space $\mathcal{H}$. The standard basis $\{|\hbar a\rangle\}_0^\infty$ will be used with the ‘principle’ quantum number $a = 0, 1, 2, \dots$ assigned by Planck’s quanta counting. Generally action amplitudes are a coherent superposition

$$|G\rangle = \sum_{a=0}^{\infty} g^a |\hbar a\rangle, \quad (1)$$

which may represent of the ground state, or “vacuum”, of some quantum system. In fact only finite, say, N elementary quantum states may be involved. Then one may restrict $CP(\infty)$ to finite dimensional $CP(N-1)$. Hereafter we will use the indices as follows: $0 \leq a \leq N$, and $1 \leq i, k, m, n, s \leq N-1$

Since any ray of action amplitude has isotropy group $H = U(1) \times U(N)$, in $\mathcal{H}$ only coset transformations $G/H = SU(N)/S[U(1) \times U(N-1)] = CP(N-1)$ effectively act. Therefore the ray representation of $SU(N)$ in C^N , in particular, the embedding of H and G/H in G , is a state-dependent parametrization.

Hence, there is a diffeomorphism between the space of the rays marked by the local coordinates in the map $U_j : \{|G\rangle, |g^j\rangle \neq 0\}, j > 0$

$$\pi_{(j)}^i = \begin{cases} \frac{g^i}{g^j}, & \text{if } 1 \leq i < j \\ \frac{g^{i+1}}{g^j} & \text{if } j \leq i < N-1 \end{cases} \quad (2)$$

and the group manifold of the coset transformations $G/H = SU(N)/S[U(1) \times U(N-1)] = CP(N-1)$. This diffeomorphism is provided by the coefficient functions Φ_α^i of the local generators (see below). The choice of the map U_j means, that the comparison of quantum amplitudes refers to the amplitude with the action $\hbar j$. The breakdown of $SU(N)$ symmetry on each action amplitude to the isotropy group $H = U(1) \times U(N-1)$ contracts the full dynamics down to $CP(N-1)$. The physical interpretation of these transformations is given by the

POSTULATE 2.

We shall assume that the unitary transformations of the action amplitudes may be identified with physical fields; i.e., transformations of the form $U(\tau) = \exp(i\Omega^\alpha \hat{\lambda}_\alpha \tau)$, where the field functions Ω^α are in the adjoint representations of $SU(N)$. The coset transformation $G/H = SU(N)/S[U(1) \times U(N-1)] = CP(N-1)$ is the quantum analog of a classical force; its action is equivalent to some physically distinguishable variation of GCS in $CP(N-1)$.

The $CP(N-1)$ manifold takes the place of the “classical phase space” [10], since its points, corresponding to the GCS, are most close to classical states of motion. Two interpretations may be given for the points of $CP(N-1)$. One of them is the “Schrödinger’s lump” [8] and the second one is the analog of the Stern-Gerlach “filter’s orientations” discussed by Fivel [11]. The basic content of their physical interpretations is that one has a *macroscopic (i.e. space-time) discriminator* of two quantum states. As such, they may be used as “yes/no” states of some two-level detector. We will use the “Schrödinger’s lump” interpretation. Let us assume that GCS described by local coordinates $(\pi^1, \dots, \pi^{N-1})$ corresponds to the original lump, and the coordinates $(\pi^1 + \delta\pi^1, \dots, \pi^{N-1} + \delta\pi^{N-1})$ correspond to the lump displaced due to measurement. Such coordinates of the lump gives the a firm geometric tool for the description of quantum dynamics during interaction which may used for a measuring process.

Then the question that we now want to raise is the following: *what “classical field”, i.e. field in space-time, corresponds to the transition from the original to the displaced lump?* In other words we would like to find the measurable physical manifestation of the lump, which we shall call the “field shell”, its space-time shape and its dynamics. The lump’s perturbations will be represented by “geometric bosons” [12] whose frequencies are not a priori given, but are defined by some field equations which should be established by means of a new variation problem. Before its formulation, we wish to introduce differential geometric construction.

We will assume that all “vacua” solutions belong to a single separable *projective Hilbert space* $CP(N-1)$. The vacuum is now the stationary point of some action functional, not a solution with the minimal energy. Energy will be

associated with vector field tangent to $CP(N-1)$ giving the rate of change of the action variation with respect to the notion of Newton-Stueckelberg-Horwitz-Piron time (world time) [13]. Dynamical space-time will be built at any GCS and, particularly, at the vacuum of some “classical” problem (see below). Therefore Minkowsky space-time appears to be functionally local (state-dependent) in $CP(N-1)$ and the space-time motion dictated by the field equations connected with two infinitesimally close “vacua”. The connection between these local space-times may be physically established by the measurement given in terms of geometry of the base manifold $CP(N-1)$.

Now we are evidences of the so-called Multiverse (omnium) concept [14, 8]. We think there is only one Universe but there exists a continuum of dynamical space-times each of them related to one point of the quantum phase space $CP(N-1)$. The standard approach, identifying the Universe with a single space-time, appears to be too strong an assumption from this point of view.

3 Local dynamical variables

The state space $\mathcal{H}$ of the field configurations with finite action quanta is a stationary construction. We introduce dynamics *by the velocities of the GCS variation* representing some “elementary excitations” (quantum particles). Their dynamics is specified by the Hamiltonian, giving time variation velocities of the action quantum numbers in different directions of the tangent Hilbert space $T_{(\pi^1, \dots, \pi^{N-1})}CP(N-1)$ which takes the place of the ordinary linear quantum scheme. The rate of the action variation gives the energy of the “particles”.

The local dynamical variables correspond to internal symmetries of the GCS and their evolution should be expressed now in terms of the local coordinates π^k . The Fubini-Study metric

$$G_{ik^*} = [(1 + \sum |\pi^s|^2)\delta_{ik} - \pi^{i^*}\pi^k](1 + \sum |\pi^s|^2)^{-2} \quad (3)$$

and the affine connection

$$\Gamma_{mn}^i = \frac{1}{2}G^{ip^*}(\frac{\partial G_{mp^*}}{\partial \pi^n} + \frac{\partial G_{p^*n}}{\partial \pi^m}) = -\frac{\delta_m^i \pi^{n^*} + \delta_n^i \pi^{m^*}}{1 + \sum |\pi^s|^2} \quad (4)$$

in these coordinates will be used. Hence the internal dynamical variables and their norms should be state-dependent, i.e. local in the state space [1, 2, 3, 12]. These local dynamical variables realize a non-linear representation of the unitary global $SU(N)$ group in the Hilbert state space C^N . Namely, $N^2 - 1$ generators of $G = SU(N)$ may be divided in accordance with the Cartan decomposition: $[B, B] \in H, [B, H] \in B, [B, B] \in H$. The $(N-1)^2$ generators

$$\Phi_h^i \frac{\partial}{\partial \pi^i} + c.c. \in H, \quad 1 \leq h \leq (N-1)^2 \quad (5)$$

of the isotropy group $H = U(1) \times U(N-1)$ of the ray (Cartan sub-algebra) and $2(N-1)$ generators

$$\Phi_b^i \frac{\partial}{\partial \pi^i} + c.c. \in B, \quad 1 \leq b \leq 2(N-1) \quad (6)$$

are the coset $G/H = SU(N)/S[U(1) \times U(N-1)]$ generators realizing the breakdown of the $G = SU(N)$ symmetry of the GCS. Furthermore, the $(N-1)^2$ generators of the Cartan sub-algebra may be divided into the two sets of operators: $1 \leq c \leq N-1$ ($N-1$ is the rank of $AlgSU(N)$) Abelian operators, and $1 \leq q \leq (N-1)(N-2)$ non-Abelian operators corresponding to the non-commutative part of the Cartan sub-algebra of the isotropy (gauge) group. Here Φ_σ^i , $1 \leq \sigma \leq N^2-1$ are the coefficient functions of the generators of the non-linear $SU(N)$ realization. They give the infinitesimal shift of the i -component of the coherent state driven by the σ -component of the unitary multipole field Ω^α rotating the generators of $AlgSU(N)$ and they are defined as follows:

$$\Phi_\sigma^i = \lim_{\epsilon \rightarrow 0} \epsilon^{-1} \left\{ \frac{[\exp(i\epsilon\lambda_\sigma)]_m^i g^m}{[\exp(i\epsilon\lambda_\sigma)]_m^j g^j} - \frac{g^i}{g^j} \right\} = \lim_{\epsilon \rightarrow 0} \epsilon^{-1} \{ \pi^i(\epsilon\lambda_\sigma) - \pi^i \}, \quad (7)$$

[15]. Then the sum of the N^2-1 the energies associated with intensity of deformations of the GCS is represented by the local Hamiltonian vector field H which is linear in the partial derivatives $\frac{\partial}{\partial \pi^i} = \frac{1}{2}(\frac{\partial}{\partial \Re \pi^i} - i\frac{\partial}{\partial \Im \pi^i})$ and $\frac{\partial}{\partial \pi^{*i}} = \frac{1}{2}(\frac{\partial}{\partial \Re \pi^i} + i\frac{\partial}{\partial \Im \pi^i})$. In other words it is the tangent vector to $CP(N-1)$

$$H = T_c + T_q + V_b = \hbar\Omega^c \Phi_c^i \frac{\partial}{\partial \pi^i} + \hbar\Omega^q \Phi_q^i \frac{\partial}{\partial \pi^i} + \hbar\Omega^b \Phi_b^i \frac{\partial}{\partial \pi^i} + c.c. \quad (8)$$

The characteristic equations for the PDE $H|E\rangle = E|E\rangle$ give the parametric representations of their solutions in $CP(N-1)$. We will identify the parameter τ in these equations with a “universal time of evolution” such as the world time [13]. This time is the measure of the configuration variation, i.e. it is a measure of the distance in $CP(N-1)$ (an evolution trajectory length in the Fubini-Study metric) expressed in time units. The energy quantization will be discussed elsewhere.

4 Dynamical quantum space-time

We shall now construct a dynamical notion of space-time in terms of internal quantum amplitudes. In this way we shall arrive at an intrinsic definition of state-dependent space-time, consistently energies from the dynamics of a quantum system. In the limit of weak interaction, with essentially free motion, one can find an approximate correspondence with the classical idea of space-time.

There are two key ideas presented here:

1. A conservation law (parallel transport) of a local Hamiltonian as characterization of particles (excitations of GCS);
2. Identification of “Lorentz spin transformation matrix” [9] of the spinor and classical Lorentz transformations of an inertial frame.

Let us introduce the concept of a “dynamical space-time” as a new construction capable of detecting the coincidences of the spinor components in the formal two-level “detector” which is a part of the full quantum configuration. The realization of this “detector” is of course the free choice of an observer. It

is important only that the chosen LDV should be invariantly connected with the coherent state with respect to one of the points of the LDV spectra. Let us assume this spectra of the LDV are known.

4.1 Embedding “Hilbert (quantum) dynamics” in space-time

If we would like to have some embedding of the “Hilbert (quantum) dynamics” in space-time we should to formalize the quantum observation (or measurement of some internal dynamical variable). This diffeomorphism between rays of $CP(N-1)$ and $SU(N)$ generators mentioned above, are realized in terms of the local $SL(2, C)$ action onto the states space C^2 as follows. The basis of these spaces form two types of vectors: the normal vector $|N\rangle$ to the tangent space at some point of $CP(N-1)$ corresponding to eigenvalue λ_D of a measuring dynamical variable $\hat{D}$ and the tangent vector $|T\rangle$, generated by the coset generators of G/H . These describe the interaction used for the measurement process. It is important to understand that the measurement i.e. comparison of the expected spinor (α_0, β_0) and the measured spinor (α_1, β_1) pave the way to embedding Hilbert space dynamics into the local dynamical space-time. One has a two-level system (logical spin $1/2$ [12]) created by the quantum question (non-self-adjoint) projector onto one of the two states $|N\rangle, |T\rangle$. Their coherent states are given by the spinors (α, β) connected with infinitesimal $SL(2, C)$ transformations, giving rise to the variation of the space-time coordinates generated by local infinitesimal Lorentz transformations.

The LDV is a vector field defined over $CP(N-1)$ and the comparison of the LDV's at different GCS's (initial and perturbed due to interaction used for measurement) require some procedure for the *identification*. It is impossible to compare expected and measured LDV “directly” (lack of correspondence due to curvature of the $CP(N-1)$ [3]). Affine parallel transport is sufficient for this purpose. Parallel transport forms the condition for the coefficient functions of the LDV leading to nonlinear field equations in local dynamical space-time.

4.2 Differential geometry of the measuring procedure

The numerical value of some observable depends as a rule on a setup (the character of motion of laboratory, type of the measuring device, field strength, etc.). However the relationships between numerical values of dynamical variables and numerical characteristics of laboratory motion, field strength, etc., should be formulated in an invariant way, since they reflect the objective character of the physical interaction used in the measurement process. The numbers obtained due to the measurements carry information which does not exist a priori, i.e. before the measurement process.

POSTULATE 3

The invariant i.e. physically essential part of information represented by the coherent states of the “logical spin $1/2$ ” is related to the space-time structure. Such a postulate is based on the observation that locally space-time is the

Lorentz-invariant manifold of points modeling different physical systems corresponding to events depleted of all physical characteristics. In principle arbitrary local coordinates may be attributed to these points. The spinor structure of the Lorentz transformations represents the transformations of the coherent states of the "logical spin $1/2$ ". Thereby we can assume the measurement of the quantum dynamical variables expressed by the spinor "creates" the local space-time coordinates. We will formulate non-linear field equations in this local space-time with the help of a variational principle referring to the generator of the quantum state deformation.

The internal dynamics of the quantum configuration given by the action amplitude should be somehow reflected in physical space-time. We shall solve the "inverse representation problem": to find a locally unitary representation of the dynamical group $SU(N)$ in the dynamical space-time where the induced realization of the coherence group $SU(2)$ of the spinor acts [1, 2]. Its components are subject to the "Lorentz spin matrix transformations" [9]. We then build the local spinor basis invariantly related to the generalized coherent state in the $CP(N-1)$ manifold. First of all we have to have the local reference frame as some type of "representation" of $SU(N)$. Each local reference frame and, hence, $SU(N)$ "representation", may be marked by the local coordinates (2). Now we should almost literally repeat differential geometry of a smooth manifold embedded in flat ambient Hilbert space $\mathcal{H} = C^N$. The geometry of this smooth manifold is the projective Hilbert space equipped with the Fubini-Study metric (3) and with the affine connection (4).

The contact between this abstract formulation of the quantum dynamics and ordinary (linear) expressions for some dynamical variable $\hat{D}$ in $\mathcal{H} = C^\infty = l_2$ may be established due to following formulas:

$$\hat{D} = \sum_{a,b \geq 0} \langle a | \hat{D} | b \rangle \hat{P}_{ab} = \sum_{a,b \geq 0} D_{ab} \hat{P}_{ab} = \sum_{a,b \geq 0} F_D^\alpha \hat{\lambda}_{\alpha,(ab)} \hat{P}_{ab}, \quad (9)$$

where projector

$$\hat{P}_{ab} = |a \rangle \langle b| = \frac{1}{\sqrt{a!b!}} : (\eta^+)^a \exp(-\eta^+ \eta) (\eta)^a :, \quad (10)$$

where $: \dots :$ means the the normal ordering of operators, and functions F_D^α obey some field equations which will be discussed later. In particular, the Hamiltonian has a similar representation

$$\hat{H} = \sum_{a,b \geq 0} \langle a | \hat{H} | b \rangle \hat{P}_{ab} = \sum_{a,b \geq 0} H_{ab} \hat{P}_{ab} = \hbar \sum_{a,b \geq 0} \Omega^\alpha \hat{\lambda}_{\alpha,(ab)} \hat{P}_{ab}, \quad (11)$$

i.e $F_H^\alpha = \hbar \Omega^\alpha$ [20]. In order to express the measurement of the "particle's field" in geometrically intrinsic terms, we assume that the GCS is expressed in local coordinates is

$$|G(\pi^1, \dots, \pi^{N-1}) \rangle = \sum_0^{N-1} g^a(\pi) |a \rangle, \quad (12)$$

where

$$g^0(\pi^1, \dots, \pi^{N-1}) = \frac{R^2}{\sqrt{R^2 + \sum_{s=1}^{N-1} |\pi^s|^2}}, \quad (13)$$

and for $1 \leq i \leq N-1$ one has

$$g^i(\pi^1, \dots, \pi^{N-1}) = \frac{R\pi^i}{\sqrt{R^2 + \sum_{s=1}^{N-1} |\pi^s|^2}}, \quad (14)$$

i.e. $CP(N-1)$ is embedded in the Hilbert space $\mathcal{H} = C^N$.

Then the rate of the ground state evolution in world time is the tangent vector $|H\rangle$ to the evolution curve $\pi^i = \pi^i(\tau)$ given by the formula

$$|H\rangle = \frac{d|G\rangle}{d\tau} = \frac{\partial g^a}{\partial \pi^i} \frac{d\pi^i}{d\tau} |a\hbar\rangle = |T_i\rangle \frac{d\pi^i}{d\tau} = H^i |T_i\rangle, \quad (15)$$

where

$$|T_i\rangle = \frac{\partial g^a}{\partial \pi^i} |a\hbar\rangle = T_i^a |a\hbar\rangle. \quad (16)$$

Then the “acceleration” is

$$|A\rangle = \frac{d^2|G\rangle}{d\tau^2} = |g_{ik}\rangle \frac{d\pi^i}{d\tau} \frac{d\pi^k}{d\tau} + |T_i\rangle \frac{d^2\pi^i}{d\tau^2} = |N_{ik}\rangle \frac{d\pi^i}{d\tau} \frac{d\pi^k}{d\tau} + \left(\frac{d^2\pi^s}{d\tau^2} + \Gamma_{ik}^s \frac{d\pi^i}{d\tau} \frac{d\pi^k}{d\tau}\right) |T_s\rangle, \quad (17)$$

where

$$|g_{ik}\rangle = \frac{\partial^2 g^a}{\partial \pi^i \partial \pi^k} |a\hbar\rangle = |N_{ik}\rangle + \Gamma_{ik}^s |T_s\rangle \quad (18)$$

and state

$$|N\rangle = N^a |a\hbar\rangle = \left(\frac{\partial^2 g^a}{\partial \pi^i \partial \pi^k} - \Gamma_{ik}^s \frac{\partial g^a}{\partial \pi^s}\right) \frac{d\pi^i}{d\tau} \frac{d\pi^k}{d\tau} |a\hbar\rangle \quad (19)$$

is normal to the “hypersurface” of the ground states. Then the minimization of this “acceleration” under the transition from point τ to $\tau + d\tau$ may be achieved by the annihilation of the tangential component

$$\left(\frac{d^2\pi^s}{d\tau^2} + \Gamma_{ik}^s \frac{d\pi^i}{d\tau} \frac{d\pi^k}{d\tau}\right) |T_s\rangle = 0 \quad (20)$$

i.e. under the condition of affine parallel transport of the Hamiltonian vector field

$$dH^s + \Gamma_{ik}^s H^i d\pi^k = 0. \quad (21)$$

We use the Gauss-Codazzi equations

$$\begin{aligned}\frac{\partial N^a}{\partial \pi^i} &= B_i^s T_s^a \\ \frac{\partial T_k^a}{\partial \pi^i} - \Gamma_{ik}^s T_s^a &= B_{ik} N^a\end{aligned}\quad (22)$$

as a condition of integrability. These give us the dynamics of the vacuum (normal) vector and the tangent vectors, i.e. one has the local reference frame dynamics modeling the “moving representation”

$$\begin{aligned}\frac{dN^a}{d\tau} &= \frac{\partial N^a}{\partial \pi^i} \frac{d\pi^i}{d\tau} + c.c. = B_i^s T_s^a \frac{d\pi^i}{d\tau} + c.c. = B_i^s T_s^a H^i + c.c.; \\ \frac{dT_k^a}{d\tau} &= \frac{\partial T_k^a}{\partial \pi^i} \frac{d\pi^i}{d\tau} + c.c. = (B_{ik} N^a + \Gamma_{ik}^s T_s^a) \frac{d\pi^i}{d\tau} + c.c. \\ &= (B_{ik} N^a + \Gamma_{ik}^s T_s^a) H^i + c.c.\end{aligned}\quad (23)$$

Note, that $0 \leq a \leq N$, but $1 \leq i, k, m, n, s \leq N-1$. The tensor B_{ik} of the second quadratic form of the ground states “hypersurface” is as follows:

$$B_{ik} = \langle N | \frac{\partial^2}{\partial \pi^i \partial \pi^k} | G \rangle. \quad (24)$$

Now one should build the spinor in the local basis $(|N\rangle, |D\rangle)$ for the quantum question with respect to the measurement of some local dynamical variable $\vec{D}$. We will assume that there is *natural state* $|\tilde{D}\rangle$ of the quantum system in the local reference frame representation equal to the lifting of LDV $\vec{D} \in T_\pi CP(N-1)$ into the Hilbert space $\mathcal{H}$, and there is *expected state* $|D_{expect}\rangle = \alpha_0 |N\rangle + \beta_0 |\tilde{D}\rangle$, associated with the tuning of a “measuring device”. This measuring device is associated with the local projector along the normal $|N\rangle$ onto the natural state $|\tilde{D}\rangle$. In fact it defines the covariant derivative in $CP(N-1)$. The lift-vectors $|N\rangle, |D\rangle$ are given by the solutions of (23) arising under interaction used for the measurement of the LDV $\vec{D}$. In general $|D\rangle$ is not a tangent vector to $CP(N-1)$. But the normalized vector defined as the covariant derivative $|\tilde{D}\rangle = |D\rangle - \langle Norm | D \rangle |Norm\rangle$ is a tangent vector to $CP(N-1)$ (it is convenient to take $|Norm\rangle = \frac{|N\rangle}{\sqrt{\langle N | N \rangle}}$). The operation of the $|\tilde{D}\rangle$ normalization is a projector. Indeed,

$$\begin{aligned}\widetilde{|\tilde{D}\rangle} &= |D\rangle - \langle Norm | D \rangle |Norm\rangle \\ &= |D\rangle - \langle Norm | D \rangle |Norm\rangle \\ &- \langle Norm | (|D\rangle - \langle Norm | D \rangle |Norm\rangle) |Norm\rangle \\ &= |D\rangle - \langle Norm | D \rangle |Norm\rangle = |\tilde{D}\rangle.\end{aligned}\quad (25)$$

Then at the point $(\pi^1, \dots, \pi^{N-1})$ one has two components of the spinor

$$\alpha(\pi^1, \dots, \pi^{N-1}) = \frac{\langle N | D_{expect} \rangle}{\langle N | N \rangle}$$

$$\beta_{(\pi^1, \dots, \pi^{N-1})} = \frac{\langle \tilde{D} | D_{expect} \rangle}{\langle \tilde{D} | \tilde{D} \rangle} \quad (26)$$

then at an infinitesimally close point $(\pi^1 + \delta^1, \dots, \pi^{N-1} + \delta^{N-1})$ one has the new spinor

$$\begin{aligned} \alpha_{(\pi^1 + \delta^1, \dots, \pi^{N-1} + \delta^{N-1})} &= \frac{\langle N' | D_{expect} \rangle}{\langle N' | N' \rangle} \\ \beta_{(\pi^1 + \delta^1, \dots, \pi^{N-1} + \delta^{N-1})} &= \frac{\langle \tilde{D}' | D_{expect} \rangle}{\langle \tilde{D}' | \tilde{D}' \rangle} \end{aligned} \quad (27)$$

where the basis $(|N' \rangle, |\tilde{D}' \rangle)$ is the lift of the parallel transported $(|N \rangle, |\tilde{D} \rangle)$ from the infinitesimally close $(\pi^1 + \delta^1, \dots, \pi^{N-1} + \delta^{N-1})$ back to $(\pi^1, \dots, \pi^{N-1})$.

These two infinitesimally close spinors being expressed as functions of θ, ϕ, ψ, R and $\theta + \epsilon_1, \phi + \epsilon_2, \psi + \epsilon_3, R + \epsilon_4$, may be represented as follows

$$\eta = R \begin{pmatrix} \cos \frac{\theta}{2} (\cos \frac{\phi_1 - \psi_1}{2} - i \sin \frac{\phi_1 - \psi_1}{2}) \\ \sin \frac{\theta}{2} (\cos \frac{\phi_1 + \psi_1}{2} + i \sin \frac{\phi_1 + \psi_1}{2}) \end{pmatrix} = R \begin{pmatrix} C(c - is) \\ S(c_1 + is_1) \end{pmatrix} \quad (28)$$

and

$$\begin{aligned} \eta + \delta\eta &= R \begin{pmatrix} C(c - is) \\ S(c_1 + is_1) \end{pmatrix} \\ &+ R \begin{pmatrix} S(is - c)\epsilon_1 - C(s + ic)\epsilon_2 + C(s + ic)\epsilon_3 + C(c - is)\frac{\epsilon_4}{R} \\ C(c_1 + is_1)\epsilon_1 + S(ic_1 - s_1)\epsilon_2 - S(s_1 - ic_1)\epsilon_3 + S(c_1 + is_1)\frac{\epsilon_4}{R} \end{pmatrix}. \end{aligned} \quad (29)$$

They may be connected with an infinitesimal ‘‘Lorentz spin transformations matrix’’ [9]

$$L = \begin{pmatrix} 1 - \frac{i}{2}\tau(\omega_3 + ia_3) & -\frac{i}{2}\tau(\omega_1 + ia_1 - i(\omega_2 + ia_2)) \\ -\frac{i}{2}\tau(\omega_1 + ia_1 + i(\omega_2 + ia_2)) & 1 - \frac{i}{2}\tau(-\omega_3 - ia_3) \end{pmatrix}. \quad (30)$$

Then accelerations a_1, a_2, a_3 and angle velocities $\omega_1, \omega_2, \omega_3$ may be found in the linear approximation from the equation

$$\eta + \delta\eta = L\eta \quad (31)$$

as functions of the spinor components depending on local coordinates $(\pi^1, \dots, \pi^{N-1})$. Hence infinitesimal Lorentz transformations define small ‘‘space-time’’ shifts. It is convenient to take Lorentz transformations in the following form $ct' = ct + (\vec{x}\vec{a})d\tau$, $\vec{x}' = \vec{x} + ct\vec{a}d\tau + (\vec{\omega} \times \vec{x})d\tau$, where we put $\vec{a} = (a_1/c, a_2/c, a_3/c)$, $\vec{\omega} = (\omega_1, \omega_2, \omega_3)$ [9] in order to have for τ the physical dimension of time. The coordinates x^μ of points in this space-time serve in fact for the parametrization of deformations of a ‘‘field shell’’ arising under its motion according to non-linear field equations [1, 2].

5 Field equations in the dynamical space-time

In order to find the “field shell” of the perturbed GCS one should establish some wave equations in the dynamical space-time. All these notions require more precise definitions. Namely, say, in the simplest case of $CP(1)$, the “field shell” represented in the spherical coordinates is the classical vector field $\Omega^\alpha = \frac{x^\alpha}{r}(\omega + i\gamma)$, $1 \leq \alpha \leq 3$ giving the rate of the GCS variations. The tensor fields $1 \leq \alpha \leq 8, 15, \dots, N^2 - 1$ will be discussed elsewhere. Note, that quanta numbers are now strongly connected with the tensor character of the GCS driving field Ω^α . These fields are “classical” since they are not subject to quantization directly, i.e. by the attribution of the fermionic or bosonic commutation relations. They obey nonlinear field equations having soliton-like solutions. Their internal dynamical variables like spin, charge, etc., are a consequence of their dynamical structure.

A “particle” which may be associated with the “field shell” in the dynamical space-time (see below), is now described locally by the Hamiltonian vector field H . At each point $(\pi^1, \dots, \pi^{N-1})$ of the $CP(N-1)$ one has an “expectation value” of the H defined by a measuring device. But a displaced GCS may be reached along one of the continuum paths. Therefore the comparison of two vector fields and their “expectation values” at neighboring points requires some natural rule. The comparison makes sense only for the same “particle” or for its “field shell” along some path. For this reason one should have an identification procedure. The affine parallel transport in $CP(N-1)$ of vector fields is a natural and the simplest rule for the comparison of corresponding “field shells”. Physically the identification of “particle” literally means that its Hamiltonian vector field is a Fubini-Study covariant constant.

Since we have only the unitary fields Ω^α as parameters of the GCS transformations we assume that in accordance with an equivalence-like principle under the infinitesimal shift of the unitary field $\delta\Omega^\alpha$ in the dynamical space-time, the shifted Hamiltonian field should coincide with the infinitesimal shift of the tangent Hamiltonian field generated by the parallel transport in $CP(N-1)$ during world time $\delta\tau$ [13]. Thus one has

$$\hbar(\Omega^\alpha + \delta\Omega^\alpha)\Phi_\alpha^k = \hbar\Omega^\alpha(\Phi_\alpha^k - \Gamma_{mn}^k \Phi_\alpha^m V^n \delta\tau) \quad (32)$$

and, hence, in accordance with the sufficiency criterion for the equation with non-trivial solution

$$\hbar(\delta\Omega^\alpha \delta_m^k + \Omega^\alpha \Gamma_{mn}^k V^n \delta\tau)\Phi_\alpha^m = 0 \quad (33)$$

one has following equations

$$\frac{\delta\Omega^\alpha}{\delta\tau} = -\Omega^\alpha \Gamma_{mn}^m V^n, \quad (34)$$

(there is no summation in m). We introduce the dynamical space-time coordinates x^μ as state-dependent quantities, transforming in accordance with the local Lorentz transformations $x^\mu + \delta x^\mu = (\delta_\nu^\mu + \Lambda_\nu^\mu \delta\tau)x^\nu$. The parameters of

$\Lambda_\nu^\mu(\pi^1, \dots, \pi^{N-1})$ depend on the local transformations of local reference frame in $CP(N-1)$ described in

the previous paragraph. Assuming a spherically symmetrical solution, we will use the coordinates ($x^0 = ct, x^1 = r \sin \Theta \cos \Phi, x^2 = r \sin \Theta \sin \Phi, x^3 = r \cos \Theta$). In the case of spherical symmetry, $\Omega^1 = (\omega + i\gamma) \sin \Theta \cos \Phi, \Omega^2 = (\omega + i\gamma) \sin \Theta \sin \Phi, \Omega^3 = (\omega + i\gamma) \cos \Theta$ and in the general case of the separability of the angle and radial parts, one has $\Omega^\alpha = \sum C_{l,m}^\alpha Y_{l,m}(\Theta, \Phi)(\omega + i\gamma) = \sum C_{l,m}^\alpha Y_{l,m}(\Theta, \Phi)\Omega$. Then taking into account the expressions for the “4-velocity” $v^\mu = \frac{\delta x^\mu}{\delta \tau} = \Lambda_\nu^\mu(\pi^1, \dots, \pi^{N-1})x^\nu$ one has the field equation

$$v^\mu \frac{\partial \Omega}{\partial x^\mu} = -\Omega \Gamma_{mn}^m V^n, \quad (35)$$

where

$$\begin{aligned} v^0 &= (\vec{x}\vec{a}) \\ \vec{v} &= ct\vec{a} + (\vec{\omega} \times \vec{x}). \end{aligned} \quad (36)$$

or, in detail,

$$\begin{aligned} v^0 &= r(a_1 \sin \Theta \cos \Phi + a_2 \sin \Theta \sin \Phi + a_3 \cos \Theta) \\ v^1 &= cta_1 + r(\omega_2 \cos \Theta - \omega_3 \sin \Theta \sin \Phi) \\ v^2 &= cta_2 + r(\omega_3 \sin \Theta \cos \Phi - \omega_1 \cos \Theta) \\ v^3 &= cta_3 + r(\omega_1 \sin \Theta \sin \Phi - \omega_2 \sin \Theta \cos \Phi) \end{aligned} \quad (37)$$

If one wishes to find a field corresponding to a given trajectory, say, a geodesic in $CP(N-1)$, then, taking into account that any geodesic as whole belongs to some $CP(1)$, one may put $\pi^1 = e^{i\phi} \tan(\sigma\tau)$. Then $V^1 = \frac{d\pi^1}{d\tau} = \sigma \sec^2(\sigma\tau) e^{i\phi}$, and one has a linear wave equations for the gauge unitary field Ω^α in the dynamical space-time with complicated coefficient functions of the local coordinates $(\pi^1, \dots, \pi^{N-1})$. Under the assumption $\tau = wt$ this equation has following solution

$$= (F_1(r^2 - c^2t^2) + iF_2(r^2 - c^2t^2)) \exp(-2wc \int_0^t dp \frac{\tan(wp)}{A\sqrt{c^2(p^2 - t^2) + r^2}}), \quad (38)$$

where F_1, F_2 are an arbitrary functions of the interval $s^2 = r^2 - c^2t^2$, $(\vec{a}, \vec{x}) = Ar \cos(\chi)$, $A = \sqrt{a_x^2 + a_y^2 + a_z^2}$ and $r = \sqrt{x^2 + y^2 + z^2}$. The angle χ in fact is defined by a solution of the equation (4.20). We used $\chi = \pi$ since for us it is now interesting only (due to spherical symmetry) a “radial boost toward the center of the field shell”.

The general factor demonstrates the spreading of the light cone due to the boosts. Thus our results are consistent with the so-called “off-shell” idea of Horwitz-Piron-Stueckelberg [17, 18]. Namely, state-dependent Lorentz transformations intended to reach the coincidence of two infinitesimally close spinors lead to absolute acceleration (in the sense of deformation of quantum state). This is a measure of the deformation of GCS’s [16]. One may treat this as an effective self-interaction field of GCS leading to the spreading of the mass shell. This lump might serve as non-singular source of the field in QFT.




Figure 1: The factor $\exp(-2(r - \sqrt{r^2 - t^2}))$ in the vicinity of the light-cone

It is interesting to analyze the behavior of the general factor in the vicinity of the light-cone and in the remote areas. For this purpose it is convenient to put $w = 1, A = 1, c = 1$ and, using new variable $q = p/t$ one has

$$I(t, r) = \int_0^t dp \frac{\tan(p)}{\sqrt{p^2 - t^2 + r^2}} = \int_0^1 dq \frac{\tan(tq)}{\sqrt{\frac{r^2}{t^2} - 1 + q^2}}. \quad (39)$$

Let us put $a^2 = \frac{r^2}{t^2} - 1$. In the vicinity of the light-cone $a^2 \ll 1$ the vicinity of $q = 0$ gives the main contribution to the integral and, hence, $\tan(tq) \simeq tq$, therefore

$$I(t, r) \simeq I_0(t, r) = \int_0^1 dq \frac{tq}{\sqrt{a^2 + q^2}} = r - \sqrt{r^2 - t^2}. \quad (40)$$

The behavior of the factor $\exp(-2(r - \sqrt{r^2 - t^2}))$ is depicted in the Fig.1.

On the other hand, if $a^2 \gg 1$, then

$$I(t, r) \simeq I_r(t, r) = \frac{1}{a} \int_0^1 \tan(tq) dq = \frac{\tan(t)^2}{\sqrt{r^2 - t^2}}, \quad (41)$$

and the behavior of the factor $\exp(-2\frac{\tan(t)^2}{\sqrt{r^2 - t^2}})$ is depicted in the Fig.2. It is seen that for small t, r the behavior of these factors is similar.

Let us discuss now the self-consistent problem

$$v^\mu \frac{\partial \Omega^\alpha}{\partial x^\mu} = -(\Gamma_{mn}^m \Phi_\beta^n + \frac{\partial \Phi_\beta^n}{\partial \pi^n}) \Omega^\alpha \Omega^\beta, \quad \frac{d\pi^k}{d\tau} = \Phi_\beta^k \Omega^\beta \quad (42)$$

arising under the condition of parallel transport

$$\frac{\delta H^k}{\delta \tau} = \hbar \frac{\delta(\Phi_\alpha^k \Omega^\alpha)}{\delta \tau} = 0 \quad (43)$$

of the Hamiltonian field (21), [19, 20]. Again we will assume the simplest case of $CP(1)$ dynamics when $1 \leq \alpha, \beta \leq 3, \quad i, k, n = 1$. This system being split




Figure 2: The factor $\exp(-2\frac{\tan(t)^2}{\sqrt{r^2-t^2}})$ in the remote areas from the light-cone

into the real and imaginary parts takes the form

$$\begin{aligned} (r/c)\omega_t + ct\omega_r &= -2\omega\gamma F(u, v), \\ (r/c)\gamma_t + ct\gamma_r &= (\omega^2 - \gamma^2)F(u, v), \\ u_t &= \beta U(u, v, \omega, \gamma), \\ v_t &= \beta V(u, v, \omega, \gamma), \end{aligned} \quad (44)$$

where

$$\begin{aligned} \pi(t, r) &= u(t, r) + iv(t, r), \\ F(u, v) &= (2 \sin \Theta \cos \Phi u(t, r) + 2 \sin \Theta \sin \Phi v(t, r) + \\ &\quad \cos \Theta (1 - u(t, r)^2 - v(t, r)^2)) \frac{r \cos \Theta}{(1 + u(t, r)^2 + v(t, r)^2)(\bar{x}\bar{a})}, \\ U(u, v, \omega, \gamma) &= (1/2) \sin \Theta \cos \Phi \gamma (-1 + u(t, r)^2 - v(t, r)^2) + \\ &\quad \omega (\sin \Theta \cos \Phi v(t, r) u(t, r) - (1/2) \sin \Theta \sin \Phi (1 - u(t, r)^2 + v(t, r)^2)) + \cdot \quad (45) \\ &\quad \gamma (\sin \Theta \sin \Phi v(t, r) u(t, r) + \cos \Theta u(t, r)) + \omega \cos \Theta v(t, r), \\ V(u, v, \omega, \gamma) &= (1/2) \sin \Theta \cos \Phi \omega (1 - u(t, r)^2 + v(t, r)^2) + \\ &\quad \gamma (\sin \Theta \cos \Phi v(t, r) u(t, r) - (1/2) \sin \Theta \sin \Phi (1 + u(t, r)^2 - v(t, r)^2)) - \\ &\quad \omega (\sin \Theta \sin \Phi v(t, r) u(t, r) + \cos \Theta u(t, r)) + \gamma \cos \Theta v(t, r) \end{aligned}$$

It is impossible of course to solve this self-consistent problem analytically even in this simplest case of the two state system, but it is reasonable to develop a numerical approximation in the vicinity of the following exact solution. Let us put $\omega = \rho \cos \psi$, $\gamma = \rho \sin \psi$, then, assuming for simplicity that $\omega^2 + \gamma^2 = \rho^2 = \text{constant}$, the two first PDE's may be rewritten as follows:

$$\frac{r}{c} \psi_t + ct \psi_r = F(u, v) \rho \cos \psi. \quad (46)$$

The exact solution of this quasi-linear PDE is

$$\psi_{\text{exact}}(t, r) = \arctan \frac{\exp(2c\rho F(u, v)f(r^2 - c^2t^2))(ct + r)^{2F(u, v)} - 1}{\exp(2c\rho F(u, v)f(r^2 - c^2t^2))(ct + r)^{2F(u, v)} + 1}, \quad (47)$$

where $f(r^2 - c^2t^2)$ is an arbitrary function of the interval. It is useful to see the sketch of the spatial behavior of this solution. We have put $c = \rho = F(u, v) =$




Figure 3: The real part $\omega(t, r)$ of the amplitude




Figure 4: The real part $\frac{x}{r}\omega(t=1, r)$ of the spatial field

1, $f(r^2 - c^2t^2) = -(r^2 - c^2t^2)^2$. The 3D graphics of amplitudes $\omega(t, r), \gamma(t, r)$ and corresponding spatial fields are depicted in Fig 3-6. The real part is an isotropic vector field; somewhat analogous to a charge, whereas the imaginary part is an isotropic but that looks like a bubble in the force field. The physical interpretation of this solution is still an open question. Nevertheless, it is possible to give some general interpretation which, of course, requires careful investigation. It is known that quantum measurement induces some topologically non-trivial monopole-like vector potential associated with a Berry geometric phase [21]. The question is: is it possible to find a realistic fundamental gauge potential if one uses not artificial parameter space, but an inherently related projective Hilbert space $CP(N-1)$. If we treat the acceleration parameter $A = 1$ as an interaction constant, then by (45) one may interpret $S(t, r)$ as some relativistic source of a force field.




Figure 5: The imaginary part $\gamma(t, r)$ of the amplitude




Figure 6: The imaginary part $\frac{x^\alpha}{r}\gamma(t=1, r)$ of the spatial field

6 Discussion

It is clear that the *coincidence* of the pointer with some number on the scale is in fact the coincidence of the two space-time points. But one has in the quantum area literally neither pointer nor the scale; instead we deal with a “field shell”. In Einstein’s original thinking [22], he had intuitively clear classical measuring devices (clocks, scales, rods, etc.) and the intuitively clear space-time coincidence of two “points”. Without these ingredients it is difficult to image classical measurement processes and even the notion of space-time itself. However, in fact *space-time separation is state-dependent* [23, 24]. In such a situation, one should decide on what criterion of identity is physically acceptable. It has been formulated here as the affine parallel transport of the local Hamiltonian vector field over $CP(N-1)$.

A review of the proposed scheme is as follows:

- a). We discuss the representation of the $G = SU(N)$ acting on the states $|S\rangle \in \mathcal{H}$ in terms of local dynamical variables representation by the tangent vectors to $CP(N-1)$ (the operators of differentiation); these correspond to states of the action.
- b). Measurement is realized as a perturbation of the generalized coherent quantum state.
- c). Identification of quantum systems with help of the affine parallel transport agrees with Fubini-Study metric.
- d). A variational principle applied to the local Hamiltonian leads to quasi-linear PDE field equations for the “field shell”.
- e). The fundamental “field shell” of the local dynamical variables provides as a model of an extended particle serving for the establishment of a local state-dependent space-time structure.

In the framework of our model the projection acts continuously and locally along the $CP(N-1)$ trajectory of GCS onto the corresponding tangent space, since it is the covariant differentiation of vector fields representing LDV’s on $CP(N-1)$.

Deformation of GCS occurs during interaction involved with measurement. Let us discuss the dynamics of Schrödinger’s lump during measurement (see paragraph 30.10, [8]). This construction is a humane version of the Schrödinger’s cat. In distinction with so complicated a system as a poisoned cat, and indefinite displaced lump of matter, we would like discuss the deformation of GCS which is theoretically analyzable.

First of all we should note that the assumption that “the energy in each case is the same” may be correct only approximately, say, in the case of adiabatic “kicking” of the lump. The finite time of transition unavoidably leads to the acceleration of the lump of matter, to the deformation of its quantum state [16], and to the shift of mass-energy.

In the framework of our model, the GCS of the lump is “kicked” by the coset transformations of $SU(N)$ group. The coefficient functions of the $SU(N)$ generators obey some quasi-linear relativistic field equations in the local dynamical space-time [1, 2].

The difference of the masses of the original and the displaced lumps leads to different time-like Killing vectors (if any) in the vicinities of two lumps. This is an obstacle to write Schrödinger equations for superposed wave functions.

In the framework of our model we have state-dependent space-times arising as specific sections of the tangent fibre bundle over $CP(N-1)$.

References

- [1] P.Leifer, arXive: gr-qc/0503083.
- [2] P.Leifer, arXive: gr-qc/0505051.
- [3] P.Leifer, JETP Letters, **80**, (5) 367 (2004).
- [4] R.Cirelli, A.Maniá, L.Pizzocchero, Int.J. of Mod. Phys. A **6**, (12) 2133 (1991).
- [5] L.P.Hughston, "Geometry of Stochastic State Vector Reduction", Proc. R. Soc. Lond. **A452**, 953 (1996).
- [6] L.P.Hughston, "Geometric Aspects of Quantum Mechanics", Chapter 6 in: *Twistor Theory* (ed. S.A.Huggett), pp.55-79. Marcel Dekker.
- [7] S.L.Adler, L.P.Horwitz, arXive: quant-ph/9909026.
- [8] R.Penrose, *The Road to Reality*, Alfred A.Knopf, New-York, (2005).
- [9] C.W.Misner, K.S.Thorne, J.A.Wheeler, *Gravitation*, W.H.Freeman and Company, San Francisco, 1973.
- [10] W.-M.Zhang, D.H.Feng, Phys.Rep., (252) 1 (1995).
- [11] D.I.Fivel, Phys.Rev.Lett., **67**,(3), 285 (1991).
- [12] P.Leifer, arXive: hep-th/9712131.
- [13] L.P.Horwitz, arXive: hep-ph/9606330.
- [14] S.Weinberg, arXive: hep-th/0511037.
- [15] P.Leifer, Found. Phys. **27**, (2) 261 (1997).
- [16] P.Leifer, Found.Phys.Lett., **11**, (3) 233 (1998).
- [17] L.P.Horwitz, C.Piron, Helvetica Physica Acta, **46**, 316 (1973).
- [18] J.Frastai and L.P.Horwitz, Found. Phys. **25**, (10) 1495 (1995).
- [19] P.Leifer, Found.Phys.Lett., **18**, (2) 195 (2005).
- [20] P.Leifer, arXive: physics/0601201.

- [21] Y.Aharonov, *et al*, arXive: quant-ph/9708043.
- [22] A.Einstein, Ann. Phys. **17**, 821 (1905).
- [23] A.Einstein, Ann. Phys. **49**, 769 (1916).
- [24] Y.Aharonov, *et al*, arXive: quant-ph/9709031.